

Package: CropBreeding (via r-universe)

December 12, 2024

Type Package

Title Stability Analysis in Crop Breeding

Version 0.1.0

Maintainer Prity Kumari <psingh2506@gmail.com>

Description Provides tools for crop breeding analysis including

Genetic Coefficient of Variation (GCV), Phenotypic Coefficient of Variation (PCV), heritability, genetic advance calculations, stability analysis using the Eberhart-Russell model, two-way ANOVA for genotype-environment interactions, and Additive Main Effects and Multiplicative Interaction (AMMI) analysis. These tools are developed for crop breeding research and stability evaluation under various environmental conditions. The methods are based on established statistical and biometrical principles. Refer to Eberhart and Russell (1966) [doi:10.2135/cropsci1966.0011183X000600010011x](https://doi.org/10.2135/cropsci1966.0011183X000600010011x) for stability parameters, Fisher (1935) ``The Design of Experiments'' [ISBN:9780198522294](https://doi.org/10.2307/j.ctv3383774), Falconer (1996) ``Introduction to Quantitative Genetics'' [ISBN:9780582243026](https://doi.org/10.2307/j.ctv3383776), and Singh and Chaudhary (1985) ``Biometrical Methods in Quantitative Genetic Analysis'' [ISBN:9788122433764](https://doi.org/10.2307/j.ctv3383774) for foundational methodologies.

License MIT + file LICENSE

Encoding UTF-8

Imports stats, dplyr, metan, rlang

RoxygenNote 7.3.2

NeedsCompilation no

Author Prity Kumari [aut, cre]

Date/Publication 2024-12-11 16:40:02 UTC

Config/pak/sysreqs cmake libfontconfig1-dev libfreetype6-dev make libicu-dev libssl-dev

Repository <https://prity-ai.r-universe.dev>

RemoteUrl <https://github.com/cran/CropBreeding>

RemoteRef HEAD

RemoteSha 66b38ce9a252c88674480b33d7ffe6d1f6a9f26c

Contents

breeding_metrics	2
gxe_analysis_multiple	3
perform_ammi_single_trait	4
stability_analysis	5
Index	7

breeding_metrics *Breeding Metrics Calculation*

Description

This function calculates key breeding metrics such as genotypic variance, environmental variance, genotypic coefficient of variation (GCV), phenotypic coefficient of variation (PCV), heritability, and genetic advance (GA). These metrics are critical for assessing genotype performance and genetic potential in crop breeding experiments.

Usage

```
breeding_metrics(data, genotype_col, trait_col, replication_col)
```

Arguments

- data A data frame containing the dataset with required columns.
- genotype_col Character. Name of the genotype column.
- trait_col Character. Name of the trait column.
- replication_col Character. Name of the replication column.

Value

A list containing:

- ‘ANOVA’: The summary of the ANOVA table.
- ‘GenotypicVariance’: Genotypic variance, which measures the genetic variability among genotypes.
- ‘EnvironmentalVariance’: Environmental variance, which reflects the variability due to environmental factors.
- ‘PhenotypicVariance’: Phenotypic variance, the sum of genotypic and environmental variances.
- ‘GCV’: Genotypic coefficient of variation, expressed as a percentage of the mean.
- ‘PCV’: Phenotypic coefficient of variation, expressed as a percentage of the mean.
- ‘Heritability’: Broad-sense heritability, the proportion of phenotypic variance attributable to genetic variance.

- ‘GeneticAdvance’: Genetic advance under selection, a measure of genetic improvement.
- ‘GAPercentage’: Genetic advance as a percentage of the mean, indicating the relative genetic improvement.

References

Falconer, D. S. (1996). "Introduction to Quantitative Genetics". ISBN: 9780582243026. Singh, R. K., & Chaudhary, B. D. (1985). "Biometrical Methods in Quantitative Genetic Analysis". ISBN: 9788122433764.

Examples

```
set.seed(123)
data <- data.frame(
  Genotype = rep(c("G1", "G2", "G3"), each = 10),
  Replication = rep(1:10, times = 3),
  Trait = c(rnorm(10, 50, 5), rnorm(10, 55, 5), rnorm(10, 60, 5))
)
result <- breeding_metrics(data, "Genotype", "Trait", "Replication")
print(result)
```

gxe_analysis_multiple *Two-Way ANOVA for Genotype x Environment Interaction with Multiple Traits*

Description

This function performs a two-way ANOVA to analyze genotype and environment interactions for multiple traits, including replication effects. It provides separate ANOVA results for each specified trait in the dataset.

Usage

```
gxe_analysis_multiple(
  data,
  genotype_col,
  environment_col,
  replication_col,
  trait_cols
)
```

Arguments

- | | |
|-----------------|--|
| data | A data frame containing the dataset with required columns. |
| genotype_col | Character. Name of the genotype column. |
| environment_col | Character. Name of the environment column. |

```
replication_col
    Character. Name of the replication column.
trait_cols      A vector of trait column names to analyze.
```

Value

A list containing ANOVA results for each trait.

References

Fisher, R. A. (1935). "The Design of Experiments". ISBN: 9780198522294.

Examples

```
set.seed(123)
data <- data.frame(
  Genotype = rep(c("G1", "G2", "G3"), each = 12),
  Environment = rep(c("E1", "E2", "E3", "E4"), times = 9),
  Replication = rep(c("R1", "R2", "R3"), times = 12),
  Trait1 = c(rnorm(36, 50, 5)),
  Trait2 = c(rnorm(36, 150, 10)),
  Trait3 = c(rnorm(36, 250, 15))
)
anova_results <- gxe_analysis_multiple(
  data = data,
  genotype_col = "Genotype",
  environment_col = "Environment",
  replication_col = "Replication",
  trait_cols = c("Trait1", "Trait2", "Trait3")
)
print(anova_results$Trait1)
```

perform_ammi_single_trait

Perform AMMI Analysis for a Single Trait

Description

This function performs Additive Main Effects and Multiplicative Interaction (AMMI) analysis for a single trait to evaluate genotype x environment interactions. It generates biplots and PC1 vs. Trait visualizations without relying on predictions.

Usage

```
perform_ammi_single_trait(data, env_col, gen_col, rep_col, trait_col)
```

Arguments

<code>data</code>	A data frame containing the dataset with required columns.
<code>env_col</code>	Character. Name of the environment column.
<code>gen_col</code>	Character. Name of the genotype column.
<code>rep_col</code>	Character. Name of the replication column.
<code>trait_col</code>	Character. Name of the trait column to be analyzed.

Value

A list containing:

- ‘analysis’: The AMMI analysis results.
- ‘biplot’: The biplot (PC1 vs PC2).
- ‘pc1_plot’: The PC1 vs Trait plot.

Examples

```
set.seed(123)
data <- data.frame(
  GEN = rep(c("G1", "G2", "G3", "G4"), each = 12),
  ENV = rep(c("E1", "E2", "E3"), each = 4, times = 4),
  REP = rep(1:3, times = 16),
  Y = c(rnorm(12, 50, 5), rnorm(12, 55, 5), rnorm(12, 60, 5), rnorm(12, 65, 5))
)
results <- perform_ammi_single_trait(data, "ENV", "GEN", "REP", "Y")
```

Description

This function performs stability analysis for multiple traits across different environments using Eberhart and Russell’s regression model provided by the ‘metan’ package. It computes ANOVA tables and regression parameters for assessing genotype stability.

Usage

```
stability_analysis(
  data,
  genotype_col,
  environment_col,
  replication_col,
  trait_cols
)
```

Arguments

data A data frame containing the dataset with required columns.
 genotype_col Character. Name of the genotype column.
 environment_col
 Character. Name of the environment column.
 replication_col
 Character. Name of the replication column.
 trait_cols A vector of trait column names (response variables).

Value

A list containing results for each trait:

- ‘anova’: The ANOVA table for each trait.
- ‘regression’: Regression parameters for stability analysis.

References

Eberhart, S. A., & Russell, W. A. (1966). "Stability Parameters for Comparing Varieties". *Crop Science*, 6(1), 36–40. doi:10.2135/cropsci1966.0011183X000600010011x

Examples

```

if (!requireNamespace("metan", quietly = TRUE)) {
  install.packages("metan")
}
library(metan)

# Simulated dataset
set.seed(123)
data <- data.frame(
  Genotype = rep(c("G1", "G2", "G3"), each = 12),
  Environment = rep(c("E1", "E2", "E3", "E4"), times = 9),
  Replication = rep(1:3, times = 12),
  Trait1 = c(rnorm(36, 50, 5)),
  Trait2 = c(rnorm(36, 150, 10)),
  Trait3 = c(rnorm(36, 250, 15))
)

results <- stability_analysis(
  data = data,
  genotype_col = "Genotype",
  environment_col = "Environment",
  replication_col = "Replication",
  trait_cols = c("Trait1", "Trait2", "Trait3")
)

print(results$Trait1$anova)
print(results$Trait1$regression)

```

Index

breeding_metrics, 2
gxe_analysis_multiple, 3
perform_ammi_single_trait, 4
stability_analysis, 5